



Martial arts based meditative techniques reduce pain and emotional distress and modulate underlying cortico-limbic circuitry in children with cancer



TNP2 LAB
TAKING BRAIN SCIENCE TO A NEW HIGH

Natalie Hardy, Allesandra Iadipaolo, Cindy Cohen, Elimelech Goldberg, Jeffrey Taub, Felicity Harper, Kristopher Dulay, Rebecca Cramer, Autumm Heeter, Shelley Paulisin, Martin H. Bluth, Craig Peters, Farrah Elrahal, Christine Rabinak, Hilary A. Marusak

¹Department of Pharmacy Practice, EACPHS, WSU; ²Kids Kicking Cancer, Southfield MI; ³Department of Pediatrics, SOM, WSU; ⁴Children's Hospital of Michigan, Detroit MI; ⁵Population Studies and Disparities Research Program, Karmanos Cancer Institute, Detroit MI; ⁶Department of Oncology, SOM, WSU; ⁷Department of Pathology, SOM, WSU; ⁸Department of Pharmaceutical Sciences, EACPHS, WSU; ⁹Department of Psychiatry & Behavioral Neuroscience, SOM, WSU; ¹⁰Translational Neuroscience Graduate Program, SOM, WSU

WAYNE STATE
Eugene Applebaum
College of Pharmacy
and Health Sciences

Background

- Research shows that meditative techniques are effective for reducing pain and emotional distress in adults and more recently in children. However, compliance to meditation and other psychosocial interventions is low, particularly among children.
- Kids Kicking Cancer (KCC) is a nonprofit organization that provides free martial-arts based (MAB) meditative therapy to help children with cancer, other chronic illnesses, and their siblings cope with pain and emotional distress. Martial arts are inherently appealing to children, resulting in high compliance rates.
- Neuroimaging studies in adults show that meditation modulates activity and connectivity in cortico-limbic regions, which are involved in the subjective experience of pain and distress.
- Despite the widespread application of these techniques in pediatric samples, no neuroimaging studies have examined the effects of meditation on neurobiological mechanisms underlying pain and distress in children.



Objectives

1. To examine the effects of a brief (4-week) KCC intervention on pediatric pain and distress, and underlying cortico-limbic neural circuitry.
2. To examine the effects of (MAB) meditative and non-meditative emotion regulation techniques on underlying cortico-limbic circuitry.

Methods


- 14 youth with cancer (5-17 years, 6 females) completed 4 weekly Kids Kicking Cancer classes. Children reported pain and distress before and after each class using Likert scales. Pain and distress were also recorded for other class members, including children with other chronic illnesses (e.g., diabetes, chronic pain, sickle cell), and their healthy siblings.
- A subset of youth completed functional magnetic resonance imaging scanning before and after the Kids Kicking Cancer intervention. Functional connectivity of cortico-limbic neural circuitry was compared before and after the intervention.
- During the second scan, participants completed an fMRI task that involved viewing distress-inducing videoclips. Prior to each clip, participants received MAB meditative (attention to breath, mindful acceptance) or non-meditative instructions (distraction [count backwards from ten]) or to passively view the clip. Participants rated their distress after each clip using a Likert scale. Distress and cortico-limbic activity were compared between each instruction.

Results

MAB meditative relative to non-meditative instructions are associated with lower cortico-limbic activity

Attention to breath < Count backwards from ten



Attention to breath < Passive viewing



dACC, dorsal anterior cingulate cortex. Masked within anatomical dACC and insula regions of interest

Significant reduction in pain and emotional distress over the course of a 1-hr KCC class

Significant reduction in cortico-limbic connectivity after a 4-week KCC intervention



Conclusions

- These results provide new evidence that participation in KCC (1) lowers pediatric pain and emotional distress and (2) reduces functional connectivity in cortico-limbic brain systems critical for the perception and regulation of pain and distress. In addition, (3) active meditation, as taught in KCC, modulates the neural mechanisms underlying pain and distress.
- Importantly, the reduction in pain and distress over the course of a 1-hr KCC class was observed not only in children with cancer, but in children with other chronic illnesses (e.g., sickle cell) and their healthy siblings, suggesting that the therapy of KCC may be beneficial for a range of pediatric populations. For example, children with sickle cell experience frequent pain or sickle crises, and recent studies indicate that siblings of children with chronic diseases are at high risk of anxiety, depression, and other psychosocial problems.
- Although non-meditative techniques (e.g., distraction) have also been shown to reduce pediatric pain and distress and are widely used in pediatric healthcare settings, our data suggest that meditative techniques are more effective at modulating cortico-limbic circuitry.
- Given that meditative interventions affect the brain, meditative interventions may be longer lasting or stronger than non-meditative techniques. Future studies should test whether reduction of cortico-limbic response may help to alleviate long-term consequences of repeated pain exposures or distress (e.g., chronic pain, anxiety), and if these techniques are beneficial for reducing neural signatures of pain/distress during painful treatment-related procedures (e.g., lumbar punctures)

Acknowledgments

A very special thank you to the children.

